

$\text{Ly}\alpha$ Fluorescent Excitation of Fe II in Active Galactic Nuclei

T. A. A. Sigut

Department of Physics and Astronomy, The University of Western Ontario

and

Anil K. Pradhan

Department of Astronomy, The Ohio State University

Received _____; accepted _____

To Be Published in The Astrophysical Journal Letters

ABSTRACT

We have calculated Fe II emission line strengths for Active Galactic Nuclei Broad-Line Regions using precise radiative transfer and Iron Project atomic data. We improve the treatment of all previously considered excitation mechanisms for the Fe II emission, continuum fluorescence, collisional excitation, fluorescence by self-overlap among the iron lines, and fluorescent excitation by $\text{Ly}\alpha$. We demonstrate that $\text{Ly}\alpha$ fluorescence is of fundamental importance in determining the strength of the Fe II emission. In addition to enhancing the ultraviolet and optical Fe II flux, $\text{Ly}\alpha$ fluorescence also results in significant near-infrared Fe II emission in the $\lambda\lambda 8500 - 9500 \text{ \AA}$ wavelength range. New observations are suggested to probe this effect in strong Fe II emitting quasars.

Subject headings: quasars: emission lines — radiative transfer — line: formation

1. Introduction

One of the most puzzling instances of Fe II line formation occurs in the spectra of Active Galactic Nuclei (AGN) with Broad-Line Regions (BLR) which exhibit strong ultraviolet Fe II emission ($UV \equiv \lambda\lambda 2000 - 3000 \text{ \AA}$) and often strong optical Fe II emission ($OP \equiv \lambda\lambda 3000 - 6000 \text{ \AA}$) as well. The widths of the iron lines indicate formation in the dense BLR, yet standard photoionization calculations, successful in interpreting many other BLR features, fail to account for the strength of the Fe II emission (Wills, Netzer & Wills 1985; Joly 1993). Observed total Fe II/H β ratios are ≈ 12 , whereas standard models predict ratios less than 8, typically 3 - 5, with the largest values resulting from enhanced iron abundances. There is also a class of “superstrong” Fe II emitters with Fe II/H $\beta \approx 30$ which seem unable to be explained by traditional photoionization modeling (Joly 1993; Graham *et al* 1996).

Excitation mechanisms advanced for the Fe II emission include continuum fluorescence via the UV resonance lines, self-fluorescence via overlapping Fe II transitions, and collisional excitation, with the latter thought to contribute the bulk of the Fe II emission. However, Wills *et al* (1985) demonstrated that model BLR calculations including these excitation mechanisms cannot account for the observed strengths of the Fe II lines. Elitzur & Netzer (1985, hereafter EN) extended this calculation to include Ly α fluorescent excitation but found its contribution to be negligible. Nevertheless, Penston (1987) argued that strong indirect observational evidence for the importance of Ly α fluorescence is found in the presence of unexpected UV Fe II multiplets in the spectrum of the symbiotic star RR Tel which seem attributable only to cascades from higher levels pumped by Ly α fluorescence. The emission nebulae of symbiotics offer densities and ionization parameters similar to the BLRs of AGN. Recently, Graham *et al* (1996) have identified emission from these UV Fe II multiplets expected to be preferentially strengthened by Ly α in the spectrum of the

ultra-strong Fe II emitter 2226-3905.

In this *Letter*, we improve the modeling of all of the mentioned excitation mechanisms by using new, accurate atomic data and precise radiative transfer, and we demonstrate that fluorescent excitation by Ly α can be of prime importance. A key feature of Ly α fluorescent excitation of Fe II is significant near-IR Fe II emission in the wavelength range $\lambda\lambda 8500 - 9500$ Å. We suggest that a currently well-known, but unidentified, emission feature near $\lambda 9200$ in AGN spectra may be, at least in part, a spectral signature of this process.

2. Calculations

Fe I - IV were considered with Fe I, Fe III, and Fe IV each represented by a ground state photoionization cross section and total, unified radiative plus dielectronic recombination rate from the Iron Project calculations (see Bautista & Pradhan 1998). The Fe II atomic model consists of 262 fine-structure levels with energies from Johansson (1978; 1992 private communication). The model atom is complete in quartet and sextet levels for energies below 0.849 Rydbergs, allowing full inclusion of all Ly α fluorescently excited transitions from a⁴D. This atom is a subset of the much larger 827 level Fe II atom used by Sigut & Pradhan (1996).

Transition probabilities for Fe II were drawn from 3 sources, Fuhr, Martin & Wiese (1988), Nahar (1995), and Kurucz (1991, private communication). Over 3,400 Fe II radiative transitions were included in the non-LTE solution. Photoionization cross sections for all Fe II levels were taken from Nahar & Pradhan (1994). A level-specific radiative plus dielectronic recombination coefficient was used for each Fe II quartet and sextet level. For collisional excitation of Fe II, we adopted the R-matrix results of Zhang & Pradhan (1995) and Bautista & Pradhan (1998) and used the Gaunt-factor formula for all remaining

high-lying, dipole transitions. Charge transfer recombination rates to Fe II and III were taken from Kingdon & Ferland (1996). Charge transfer ionization of Fe II was also included as Fe III + e^- recombination is to the Fe II ground state (Neufeld & Dalgarno 1987).

The coupled equations of radiative transfer and statistical equilibrium were solved with the accelerated lambda-iteration method of Rybicki & Hummer (1992). As overlap between the Fe II transitions is complex, the full preconditioning strategy was implemented. Complete frequency redistribution (CRD) over depth-dependent Doppler profiles was assumed for all iron transitions. The calculations were performed in plane-parallel, one dimensional, continuum illuminated clouds, each computed for a given ionization parameter and total particle density, assuming constant total pressure. The run of T_e and N_e , the photoionizing radiation field, and the continuous opacities and source functions were obtained with the CLOUDY code (Ferland 1991). This cloud structure was fixed during the iron non-LTE calculation. The solar iron abundance was used, $\log(N_{\text{Fe}}/N_{\text{H}}) + 12 = 7.477$.

The Ly α source function can be written without significant error as

$$S_{\nu}^{\text{Ly}\alpha} = \left(\frac{n_{2p} A_{2p,1s}}{n_{1s} B_{1s,2p} - n_{2p} B_{2p,1s}} \right) \frac{\psi_{\nu}}{\phi_{\nu}} \quad (1)$$

because stimulated emission is not important. The quantity in brackets is the usual CRD source function. The absorption profile, ϕ_{ν} , was taken to be a depth-dependent Voigt profile with a width set by natural broadening. The emission profile, ψ_{ν} , however, depends on the radiation field (Mihalas 1978). We have retained the CLOUDY H I 1s and 2p level populations, but have explicitly computed the emission profile in the partial coherent scattering approximation as formulated by Kneer (1975). Assuming a fraction γ_c of Ly α photons are coherently scattered in the atom’s rest frame, the ratio of emission to absorption profile is

$$\frac{\psi_{\nu}}{\phi_{\nu}} = 1 + \gamma_c \left(\frac{J_{\nu} \cdot \int a_{\nu',\nu} \phi_{\nu'} d\nu' - \int a_{\nu',\nu} \phi_{\nu'} J_{\nu'} d\nu'}{\int \phi_{\nu'} J_{\nu'} d\nu'} \right), \quad (2)$$

where J_ν is the mean intensity, and $a_{\nu',\nu}$ is a symmetric function that effects the transition from CRD in the line core to coherent scattering in the wings. We have taken the form of $a_{\nu',\nu}$ from Vernazza, Avrett & Loeser (1981) and adopted $\gamma_c = 0.98$. We solved the radiative transfer problem implied by equation (2) separately with an iterative technique, and then the $\text{Ly}\alpha$ source function and opacity were incorporated into the appropriate monochromatic source functions during the solution of the radiative transfer equations for the Fe II transitions. We have verified our solution of equation (2) by matching the $\text{Ly}\alpha$ profiles tabulated for BLR clouds by Avrett & Loeser (1988) using their models and hydrogen populations.

3. Results

Figure 1 compares the predicted Fe II flux with and without $\text{Ly}\alpha$ pumping in BLR models with two different ionization parameters (U_{ion}), 10^{-2} and 10^{-3} . These values are chosen to be representative of typical Fe II-emitting clouds, but do not exhaust the range of possibilities. Both models assumed a total particle density $4 \cdot 10^9 \text{ cm}^{-3}$, a total column density of 10^{23} cm^{-2} , and zero microturbulence. The shape of the photoionizing continuum was that of Mathews & Ferland (1987).

For $U_{\text{ion}} = 10^{-2}$, the influence of $\text{Ly}\alpha$ is large, more than doubling the total emitted flux, giving $\text{Fe II}/\text{H}\beta = 2.4$ for UV emission and 0.78 for optical emission, as compared to 1.1 and 0.29, respectively, without $\text{Ly}\alpha$ (the $\text{H}\beta$ flux was taken from the CLOUDY model). The strongest flux enhancements occur as noted by Penston (1987). The most important fluorescent excitation rates are from low-lying a^4D^e levels to the $^4\text{D}^o$, $^4\text{F}^o$, $^4\text{P}^o$, $^4\text{S}^o$, and $^6\text{F}^o$ symmetries of the $3\text{d}^6(^5\text{D})5\text{p}$ configuration (Johansson & Jordan 1984). Cascades from these levels to e^4D and e^6D produce strong emission lines in the range $\lambda\lambda 8500 - 9500 \text{ \AA}$ (see Fig. 2). In turn, cascades from e^4D and e^6D to odd parity levels near 5 eV enhance

multiplets 399, 391, 380 ($\sim \lambda 2850$), 373 ($\sim \lambda 2770$), and 363 ($\sim \lambda 2530$); these lines are the unexpected UV multiplets in the RR Tel spectrum noted by Penston and the multiplets identified by Graham *et al* (1996) in 2226-3905. Penston estimated that $\approx 20\%$ of the UV Fe II flux comes out in these multiplets, which compares favorably with the $\approx 30\%$ found in the current model. Subsequent cascades then enhance the optical and UV fluxes from transitions among the lower levels. While transitions from a^4D within $\pm 3 \text{ \AA}$ of $\text{Ly}\alpha$ are the most important in the fluorescent process (see the listing of Johansson & Jordan 1984), our treatment of $\text{Ly}\alpha$ includes the entire profile within $\sim \pm 50 \text{ \AA}$. Including pumping only within $\pm 3 \text{ \AA}$ reduces the flux ratios to 2.1 for UV emission and 0.70 for optical emission for the case of $U_{\text{ion}} = 10^{-2}$.

For $U_{\text{ion}} = 10^{-3}$, the effect of $\text{Ly}\alpha$ pumping is not as large, particularly on the UV flux. The Fe II/ $\text{H}\beta$ ratio generally increases with decreasing ionization parameter (even without $\text{Ly}\alpha$ pumping) because of the larger sensitivity of the hydrogen level populations to temperature.

As noted by Penston (1987), and confirmed by the current calculation (Fig. 2), strong observational evidence for $\text{Ly}\alpha$ pumping of Fe II would come from the identification of the initial decays from the pumped 5p levels to e^4D and e^6D_4 which give rise to transitions in the $\lambda\lambda 8500 - 9500 \text{ \AA}$ wavelength range. We note that there is a prominent unidentified feature near $\lambda 9200$ in quasar spectra as discussed by Morris & Ward (1989) and Osterbrock, Shaw & Veilleux (1990). An identification as solely H I Pa9 $\lambda 9229$ would render it anomalously strong compared to Pa8 and Pa10. Morris & Ward tentatively assign this feature to Mg II $4p \text{ } ^2P_{3/2,1/2}^o - 4s \text{ } ^2S_{1/2}$ ($\lambda 9218, 9244$) as formed by cascades from $5p \text{ } ^2P_{3/2,1/2}^o$ pumped by $\text{Ly}\beta$. They discuss possible contributions from Fe II but discount this possibility based on the apparent absence of a line at $\lambda 8927$. Our calculations indicate the $\lambda 8927$ Fe II line is significantly weaker than the main feature near $\lambda 9200$ (see Fig. 2). We have

performed a preliminary non-LTE calculation for Mg II including Ly β fluorescence in a manner similar to that described for Ly α (but with $\gamma_c = 0.4$) and find that while the Mg II transitions in this region are strengthened by Ly β by nearly a factor of 5, they are still only minor contributors to the flux in the $\lambda 9200$ region ($< 5\%$). Hence, the $\lambda 9200$ feature may be, in part, a spectral signature of Ly α pumping of Fe II. As noted by Osterbrock *et al*, observations in this wavelength region are hampered by telluric H₂O absorption bands, which require careful removal, and, for available observations, falling CCD responses in the near-IR. New observations of this wavelength region are warranted.

The current efficiency of Ly α pumping is in disagreement with the EN calculation. An important difference is in the adopted oscillator strengths of the pumped transitions from a⁴D. The Kurucz (1981) values are roughly a factor of 10 smaller than those obtained by either Nahar (1995) or Kurucz (1991, private communication). Reducing the oscillator strengths for the pumped transitions within $\pm 3\text{ \AA}$ of Ly α by a factor of 10 results in a strong reduction in the effect of Ly α fluorescence, giving for the $U_{\text{ion}} = 10^{-2}$ model of Figure 1, Fe II/H β ratios of 1.28 for UV emission and 0.36 for optical emission. The near-IR emission (Fig. 2) is also substantially reduced from Fe II/H $\beta = 0.326$ to 0.061. EN also used a simple escape probability treatment of line overlap, and a simpler treatment of Ly α , so significant differences with the current work are perhaps expected. Finally, we note that our atom is essentially complete in the large number of quartet and sextet Fe II energy levels below the 3d⁶(⁵D)5p configuration, and that our branching ratios from the pumped levels are expected to be considerably more accurate than those used by EN.

4. Conclusions

Using precise radiative transfer and accurate atomic data for Fe II, we demonstrate that Ly α fluorescent excitation can more than double the Fe II flux in the UV and optical

regions. We also predict Fe II emission in the $\lambda\lambda 8500 - 9500 \text{ \AA}$ region as a direct result of Ly α fluorescence. Both of these results can be viewed as theoretical confirmation of the role of Ly α fluorescence suggested by Penston (1987) on the basis of observations. New observations of the $\lambda\lambda 8500 - 8500 \text{ \AA}$ wavelength range in strong Fe II emitters are highly desirable, and would provide valuable constraints on modeling Ly α fluorescence. Potentially new information on the Ly α profile within single BLR clouds could be obtained, something which is impossible based on observations of the highly broadened Ly α profile itself.

A detailed discussion of the variation of the Fe II emission with the physical parameters defining the BLR clouds (using a much larger 827 level Fe II atom which includes the doublet spin system), and the implications for the ‘Fe II/H β ’ problem, will be presented by Sigut & Pradhan (1998).

We are also extending our calculations to overcome some of the limitations of the present work, in particular, to incorporate all important ions into a full solution of the non-LTE radiative transfer problem for the thermal structure of AGN BLR clouds. This will enable the Ly α -Fe II problem to be solved within a self-consistent framework.

This work was supported by the US NSF through grants to the NCAR High Altitude Observatory and to AKP (PHY-9421898). This work was begun while TAAS was an NSERC postdoctoral fellow at HAO. TAAS wishes to thank J. M. Marlborough and J. D. Landstreet for support through their NSERC grants.

REFERENCES

- Avrett, E. H., & Loeser, R. 1988 ApJ 331, 211
- Bautista, M. A., & Pradhan, A. K. 1998 ApJ 492, 650
- Elitzur, M., & Netzer, H. 1985 ApJ 291, 464
- Ferland, G. J. 1991 *Hazy: An Introduction to Cloudy*, OSU Internal Report 91-01
- Fuhr, J. R., Martin, G. A., & Wiese, W. L. 1988 J. Phys. Chem. Ref. Data 17, 1
- Graham, M. J., Clowes, R. G., & Campusano, L. E. 1996 MNRAS 279, 1349
- Johansson, S., 1978 Physica Script 18, 217
- Johansson, S., Jordan, C. 1984 MNRAS 210, 239
- Joly, M. 1993 Ann. Phys. Fr. 18, 241
- Kingdon, J. B., & Ferland, G. J. 1996 ApJS 106, 205
- Kneer, F. ApJ 200, 367
- Kurucz, R. L. 1981, *Semi-empirical Calculation of gf values : Fe II*, Smithsonian Astrophysical Observatory Report 390 (Cambridge)
- Mathews, W. G., & Ferland, G. J. 1987 ApJ 323, 456
- Mihalas, D. 1978 *Stellar Atmospheres*, 2nd edition (W. H. Freeman & Company : San Fransico)
- Morris, S. L., & Ward, M. J. 1989 ApJ 340, 713
- Nahar, S. N. 1995 A&A 293, 967
- Nahar, S. N., & Pradhan A. K. 1994 J. Phys. B: At. Mol. Opt. Phys. 27, 429
- Neufeld, D. A., & Dalgarno, A. 1987 Phys. Rev. A 35, 3142
- Osterbrock, D. E., Shaw, R. A., & Veilleux, S. 1990 ApJ 352, 561

- Penston, M. V. 1987 MNRAS 229, 1p
- Rybicki, G. B., & Hummer, D. G. 1992 A&A 262, 209
- Sigut, T. A. A., & Pradhan, A. K. 1996 BAAS 28, 1405
- Sigut, T. A. A., & Pradhan, A. K. 1998 ApJ to be submitted
- Vernazza, J. E., Avrett, E. H., & Loeser, R. 1981 ApJS 45, 635
- Wills, B. J., Netzer, H., & Wills, D. 1985 ApJ 288, 94
- Zhang, H. L., & Pradhan, A. K. 1995 A&A 293, 953

Fig. 1.— The predicted Fe II spectrum. The spectra were convolved with a Gaussian of $1/e$ width of 2000 km s^{-1} . Covering fractions of the illuminated and shielded cloud faces were assumed to be 5% each. The ratios of Fe II fluxes with $\text{H}\beta$ are computed with the optical (OP) and ultraviolet (UV) wavelength regions defined in the introduction.

Fig. 2.— An enlargement of the near-IR Fe II spectrum for the $U_{\text{ion}} = 10^{-2}$ model of Figure 1. The spectra were convolved with a Gaussian of $1/e$ width of 500 km s^{-1} . ‘ $\text{Ly}\alpha$ (1/10)’ indicates that the A_{ji} values of the pumped Fe II transitions within $\pm 3 \text{ \AA}$ of $\text{Ly}\alpha$ were scaled by 0.1. Fe II(IR) refers to the flux between $0.7 - 1.0 \mu\text{m}$.



